

ON THE GENERALITY OF METHODS OF MATHEMATICAL PHYSICS AND NUMERICAL ANALYSIS FOR SOME BOUNDARY VALUE PROBLEMS-I

T. S. Vashakmadze¹, Y. F. Gülver²

¹I.Vekua Institute of Applied Mathematics and Department of
Mathematics of Iv. Javakhishvili Tbilisi State University
11 University Str., Tbilisi 0186, Georgia
tamazvashakmadze@gmail.com

²Pîrî Reis University, Engineering Faculty
Tuzla Postane Mah. Eflatun Sk. No:8, 34940 İstanbul, Türkiye
yfgulver@pirireis.edu.tr

(Received 23.07.2021; accepted 06.12.2021)

Abstract

We consider the problems of creation the convergence difference schemes and numerical realization for the estimate of order of arithmetic operations needed for finding an approximate solution of Dirichlet problem for biharmonic equation in multidimensional cube. In this paper we consider and analyse one-dim case.

Keywords and phrases: Boundary value problems, generalised numerical solution, finite difference method, variational method.

AMS subject classification (2010): 34K10, 35D30, 65N06, 65N99.

1 Introduction

This study represents an ideological breakthrough that is essentially manifested when considering multidimensional boundary value problems (BVPs). In general, there are two directions: (1) The possibility to study BVPs by applying methods of mathematical physics together with methods of numerical analysis, which requires creation of new type functionals and determination of the properties of corresponding operators; (2) The possibility of effective numerical realization of generalized solutions or relatively less smooth boundary value problems.

As a first step, to see the effect of the method, we consider very simple 1D BVPs of second order operator. The results are compared with the results of an exact solution and direct application of classical finite difference method.

2 BVP-1: Second Order ODE

Consider the following BVP

$$\begin{aligned} u'' &= f, \\ u|_{\partial\Omega} &= 0, \\ \text{with } u &:= u(x), \quad f := f(x) := 12x^2 - 12x + 2, \quad x \in \Omega := (0, 1). \end{aligned} \quad (1)$$

3 Solution of the BVP-1

The exact solution of equation (1) is

$$u_e = x^2(1 - x)^2 \quad (2)$$

The approximate solution is found by applying variational and numerical methods together. After some variational manipulations (see for example [1]) the approximate functional related to BVP-1 is derived for $i = \overline{1, n-1}$ as:

$$\int_{\Omega} [(u')^2 + 2fu] dx \cong \tilde{I}(\cdots, u_i, \cdots), \quad (3)$$

Let $\int_{\Omega} (u')^2 dx \cong \tilde{I}_L(\cdots, u_i, \cdots)$, $-\int_{\Omega} 2fudx \cong \tilde{I}_R(\cdots, u_i, \cdots)$. Then the corresponding Euler equation of BVP-1 requires that

$$\frac{\partial \tilde{I}_L}{\partial u_i} \cong \frac{\partial \tilde{I}_R}{\partial u_i}. \quad (4)$$

From now on, techniques of numerical analysis is to be applied to get algebraic system of equations from (4). Ω region can be discretised by dividing in n parts with a uniform increment of h , as $h = 1/n = x_{i+1} - x_i, (i = \overline{0, n-1})$. The left hand side integral given in (4) is derived by selecting 2-point central difference stencil for the first order derivative and covering the region Ω by passing through each inner point of the region $x_i (i = \overline{1, n-1})$. So

$$2h \sum_{i=1}^{n-1} \left(\frac{u_{i+1} - u_{i-1}}{2h} \right)^2 \cong 2 \int_0^1 (u')^2 dx - \int_0^h (u')^2 dx - \int_{(n-1)h}^{nh} (u')^2 dx. \quad (5)$$

From this relation the left hand side integral given in (4) can be derived

as

$$\begin{aligned} \int_0^1 (u')^2 dx &\cong \tilde{I}_L(\cdots, u_i, \cdots) := h \sum_{i=1}^{n-1} \left(\frac{u_{i+1} - u_{i-1}}{2h} \right)^2 + \\ &\frac{1}{2} \int_0^h (u')^2 dx + \frac{1}{2} \int_{(n-1)h}^{nh} (u')^2 dx, \end{aligned} \quad (6)$$

where the integrals near to the boundary points can be approximated by taking 3-point one sided derivative stencil for the end points and the integrals are approximated by applying closed trapezoidal rule as follows [2]

$$\begin{aligned} \int_0^h (u')^2 dx &\cong h \cdot \frac{(u'_0)^2 + (u'_1)^2}{2} \\ u'_0 &= \frac{1}{2h}(-3u_0 + 4u_1 - u_2), \\ u'_1 &= \frac{u_2 - u_0}{2h}, \\ \int_{(n-1)h}^{nh} (u')^2 dx &\cong h \cdot \frac{(u'_{n-1})^2 + (u'_n)^2}{2}, \\ u'_{n-1} &= \frac{u_n - u_{n-2}}{2h}, \\ u'_n &= \frac{1}{2h}(3u_n - 4u_{n-1} + u_{n-2}). \end{aligned} \quad (7)$$

The right hand side integral given in (4) can be calculated by the open trapezoidal rule as

$$-2 \int_0^1 f u dx \cong \tilde{I}_R(\cdots, u_i, \cdots) := -2h \sum_{i=1}^{n-1} (f u)_i \quad (8)$$

Now, as required by (4), the partial derivative of the left and right hand side approximate integrals given in (6) and (8) respectively can be taken,

the results are given below

$$\begin{aligned}
\frac{\partial \tilde{I}_L}{\partial u_i} &= \frac{1}{h} \left(u_i - \frac{1}{2}u_{i+2} - \frac{1}{2}u_{i-2} \right), \quad (i = \overline{3, n-3}), \\
\frac{\partial \tilde{I}_L}{\partial u_1} &= \frac{1}{h} \left(\frac{5}{2}u_1 - \frac{1}{2}u_2 - \frac{1}{2}u_3 \right), \\
\frac{\partial \tilde{I}_L}{\partial u_2} &= \frac{1}{h} \left(-\frac{1}{2}u_1 + \frac{5}{4}u_2 - \frac{1}{2}u_4 \right), \\
\frac{\partial \tilde{I}_L}{\partial u_{n-2}} &= \frac{1}{h} \left(-\frac{1}{2}u_{n-4} + \frac{5}{4}u_{n-2} - \frac{1}{2}u_{n-1} \right), \\
\frac{\partial \tilde{I}_L}{\partial u_{n-1}} &= \frac{1}{h} \left(-\frac{1}{2}u_{n-3} - \frac{1}{2}u_{n-2} + \frac{5}{4}u_{n-1} \right), \\
\frac{\partial \tilde{I}_R}{\partial u_i} &= -2hf_i.
\end{aligned} \tag{9}$$

Carefull consideration of the algebraic system given in (9) together with (4), reveals that the system can be divided into two subsystem for odd and even indices. Then the whole system $(n-1) \times (n-1)$ can be solved simply by reducing it to (4×4) system as follows.

Let us consider when $n = 2m$; let $u_\alpha = \nu_\alpha, u_{2m-\alpha} = w_\alpha (\alpha = 1, 2)$ are parameters and $-2h^2 f(x_i) = \varphi_i (i = \overline{0, 2m})$. Thus we have the following steps:

1. Let in (9a) the indices are odd and by using (9f), we have:

$$\begin{aligned}
u_{2i+1} &= \frac{1}{2}u_{2i-1} + \frac{1}{2}u_{2i+3} + \varphi_{2i+1} \quad (i = \overline{1, m-2}) \Rightarrow u_3 = \\
&\frac{1}{2}u_5 + \frac{1}{2}\nu_1 + \varphi_3, \quad \varphi_3 = F_3, \\
u_5 &= \frac{2}{3}u_7 + \frac{1}{3}\nu_1 + F_5, \quad F_5 = \frac{4}{3} \left(\frac{1}{2}F_3 + \varphi_5 \right), \dots, \\
u_{2k+1} &= \frac{k}{k+1}u_{2k+3} + \frac{1}{k+1}\nu_1 + F_{2k+1}, \\
F_{2k+1} &= \frac{2k}{k+1} \left(\frac{1}{2}F_{2k-1} + \varphi_{2k+1} \right).
\end{aligned} \tag{10}$$

Taking $k = m-2$ in (10a) and using (9e) immediately follows:

$$-\frac{1}{2(m-1)}\nu_1 + \frac{4m-3}{2(m-1)}w_1 - \frac{1}{2}w_2 = \phi_1, \quad \phi_1 = \frac{1}{2}F_{2m-3} + \varphi_{2m-1}. \tag{11}$$

2. Let in (9a) the indices are even and by using (9f), we analogously have:

$$\begin{aligned}
u_{2i} &= \frac{1}{2}u_{2i+2} + \frac{1}{2}u_{2i-2} + \varphi_{2i} \quad (i = \overline{2, m-2}) \Rightarrow u_4 = \\
&\frac{1}{2}u_6 + \frac{1}{2}\nu_2 + \varphi_4, \quad \varphi_4 = F_4, \\
u_6 &= \frac{2}{3}u_8 + \frac{1}{3}\nu_2 + F_6, \quad F_6 = \frac{4}{3} \left(\frac{1}{2}F_4 + \varphi_6 \right), \dots, \\
u_{2k+2} &= \frac{k}{k+1}u_{2k+4} + \frac{1}{k+1}\nu_2 + F_{2k+2}, \\
F_{2k+2} &= \frac{2k}{k+1} \left(\frac{1}{2}F_{2k} + \varphi_{2k+2} \right).
\end{aligned} \tag{12}$$

Taking $k = m - 3$ in (12a) and using (9d) we have:

$$-\frac{1}{2(m-2)}\nu_2 - \frac{1}{2}w_1 + \frac{3m-4}{4(m-2)}w_2 = \phi_2, \quad \phi_2 = \frac{1}{2}F_{2m-4} + \varphi_{2m-2}. \tag{13}$$

3. Let again in (9a) the indices are odd and by using (9f), we have:

$$\begin{aligned}
u_{2i+1} &= \frac{1}{2}u_{2i-1} + \frac{1}{2}u_{2i+3} + \varphi_{2i+1} \quad (i = \overline{1, m-2}) \Rightarrow u_{2m-3} = \\
&\frac{1}{2}u_{2m-5} + \frac{1}{2}w_1 + \varphi_{2m-3}, \quad \varphi_{2m-3} = \bar{F}_{2m-3}, \\
u_{2m-5} &= \frac{2}{3}u_{2m-7} + \frac{1}{3}w_1 + \bar{F}_{2m-5}, \quad \bar{F}_{2m-5} = \frac{4}{3} \left(\frac{1}{2}\bar{F}_{2m-3} + \varphi_{2m-5} \right), \dots, \\
u_{2m-2k-1} &= \frac{k}{k+1}u_{2m-2k-3} + \frac{1}{k+1}w_1 + \bar{F}_{2m-2k-1}, \\
\bar{F}_{2m-2k-1} &= \frac{2k}{k+1} \left(\frac{1}{2}\bar{F}_{2m-2k+1} + \varphi_{2m-2k-1} \right).
\end{aligned} \tag{14}$$

Taking $k = m - 2$ in (14a) and using (9b), we have the following relation:

$$-\frac{1}{2(m-1)}w_1 + \frac{4m-3}{2(m-1)}\nu_1 - \frac{1}{2}\nu_2 = \bar{\phi}_1, \quad \bar{\phi}_1 = \frac{1}{2}\bar{F}_3 + \varphi_1. \tag{15}$$

4. Let again in (9a) the indices are even and by using (9f), we have:

$$\begin{aligned}
u_{2i} &= \frac{1}{2}u_{2i+2} + \frac{1}{2}u_{2i-2} + \varphi_{2i} \quad (i = \overline{2, m-2}) \Rightarrow u_{2m-4} = \\
&\frac{1}{2}u_{2m-6} + \frac{1}{2}w_2 + \varphi_{2m-4}, \quad \varphi_{2m-4} = \bar{F}_{2m-4}, \quad u_{2m-6} = \frac{2}{3}u_{2m-8} + \\
&\frac{1}{3}w_2 + \bar{F}_{2m-6}, \quad \bar{F}_{2m-6} = \frac{4}{3} \left(\frac{1}{2}\bar{F}_{2m-4} + \varphi_{2m-6} \right), \dots, \\
u_{2m-2k-2} &= \frac{k}{k+1}u_{2m-2k-4} + \frac{1}{k+1}w_2 + \bar{F}_{2m-2k-2}, \\
\bar{F}_{2m-2k-2} &= \frac{2k}{k+1} \left(\frac{1}{2}\bar{F}_{2m-2k} + \varphi_{2m-2k-2} \right).
\end{aligned} \tag{16}$$

Taking $k = m-3$ in (16a) and using (9c), we have the following relation:

$$-\frac{1}{2(m-2)}w_2 - \frac{1}{2}\nu_1 + \frac{3m-4}{4(m-2)}\nu_2 = \bar{\phi}_2, \quad \bar{\phi}_2 = \frac{1}{2}\bar{F}_4 + \varphi_2. \tag{17}$$

5. Finally, from the equations (11a), (13a), (15a) and (17a) we get the following reduced (4×4) system for the parameters ν_α, w_α :

$$\begin{bmatrix} a & 0 & b & c \\ 0 & d & e & b \\ b & e & d & 0 \\ c & b & 0 & a \end{bmatrix} \begin{bmatrix} \nu_1 \\ \nu_2 \\ w_2 \\ w_1 \end{bmatrix} = \begin{bmatrix} \phi_1 \\ \phi_2 \\ \bar{\phi}_2 \\ \bar{\phi}_1 \end{bmatrix} \tag{18}$$

where $a = -\frac{1}{2(m-1)}$, $b = -\frac{1}{2}$, $c = \frac{4m-3}{2(m-1)}$, $d = -\frac{1}{2(m-2)}$, $e = \frac{3m-4}{4(m-2)}$

Note that the parameters ν_α, w_α in (18) is ordered in such a way that the coefficient matrix is to be symmetric with respect to both diagonals. The determinant of this system $\Delta = \frac{5m(5m-4)}{16(m-1)(m-2)} \neq 0$.

6. The solution of the system (18) is given below in a successive order:

$$\begin{aligned}
w_1 &= \frac{\beta\rho_2 - \alpha\rho_1}{\beta\gamma - \alpha^2}, \quad w_2 = \frac{1}{\alpha}(\rho_2 - \gamma w_1), \\
\nu_1 &= \frac{1}{\alpha}(\phi_1 - bw_2 - cw_1), \quad \nu_2 = \frac{1}{d}(\phi_2 - ew_2 - bw_1),
\end{aligned} \tag{19}$$

where, $\alpha = b(cd + ae)$, $\beta = -ad^2 + ae^2 + db^2$, $\gamma = -ad^2 + ab^2 + dc^2$, $\rho_1 = -ad\bar{\phi}_2 + bd\phi_1 + ae\phi_2$, $\rho_2 = -ad\bar{\phi}_1 + cd\phi_1 + ab\phi_2$.

7. To find the remaining values $u_i (i = \overline{3, 2m-3})$, two point recurrence relations given in either (10) and (12) or in (14) and (16) can be used.

8. It is right the following theorem on the stability of the above recurrence relations:

Theorem-1. Let the right parts of the recurrence relations have the following forms:

$$F_{2k+1} = \frac{2k}{k+1} \left[\frac{1}{2} F_{2k-1} + \varphi_{2k+1} \right], \quad (k = \overline{2, m-2})$$

or

$$F_{2k+2} = \frac{2k}{k+1} \left[\frac{1}{2} F_{2k} + \varphi_{2k+2} \right], \quad (k = \overline{2, m-3}), \quad n = 2m$$

Then $|F_i| \leq \frac{1}{2m} M$, $M = \max_i |-4f(x_i)|$.

These estimates are also right for $|\bar{F}_i|$, $(i = \overline{3, 2m-3})$.

9. The numerical results of the solution of BVP-1 is shown comparatively in Table-1 for $n = 32$. The results of direct application of finite difference method (with 3-point central difference stencil approximation for the second order derivative) is shown as $DIRECT_{FD}$ and the results of combination of variational and numerical methods named as VASHA. Both results are compared with the exact result and percent relative errors are also tabulated. Some columns in the table filled with blue indicating that at those points the combined method gives the better results.

Table-1 Numerical Results for the BVP-1

Results for n=32	MIN % ERROR	MAX % ERROR	u1	u2	u3	u4	u5	u6	u7	u8
% error VASHA			16.09	13.95	0.92	2.34	1.44	0.47	1.37	0.06
u_VASHA	0.06 u8	16.09 u1	0.00076904	0.0029541	0.0072845	0.01168	0.01763	0.02310	0.02961	0.03518
u_EXACT			0.00091648	0.0034332	0.0072184	0.01196	0.01738	0.02321	0.02921	0.03516
u_DIRECT_FD	0.38 u16	3.23 u1	0.00094604	0.0034904	0.0073013	0.01207	0.01751	0.02336	0.02937	0.03534
% error D_FD			3.23	1.67	1.15	0.92	0.75	0.65	0.55	0.51
Results for n=32	u9	u10	u11	u12	u13	u14	u15	u16	BVP-1 $u'' = 12x^2 - 12x + 2$, $u _{\partial\Omega} = 0, x \in \Omega = (0,1)$. SOLN-1 $u_{\text{exact}} = x^2(1-x)^2$ Note that the remaining values are symmetric with respect to value of u16.	
% error VASHA	1.30	0.28	1.20	0.38	1.17	0.43	1.14	0.43		
u_VASHA	0.04139	0.04629	0.05150	0.05514	0.05886	0.06082	0.06272	0.06277		
u_EXACT	0.04086	0.04616	0.05089	0.05493	0.05818	0.06056	0.06201	0.06250		
u_DIRECT_FD	0.04106	0.04637	0.05111	0.05516	0.05842	0.06080	0.06226	0.06274		
% error D_FD	0.49	0.45	0.43	0.42	0.41	0.40	0.40	0.38		

It should be noted that combined application of variational and numerical methods reduces the order of derivative to be approximated to half. By this way, with this combined method, relatively less smooth boundary value problems like BVP-2 given below can be solved and generalised solutions can also be obtained.

4 BVP-2: Example for Less Smooth BVP

Consider the following BVP

$$u'' = \begin{cases} 0, & x \in \left[0, \frac{1}{3}\right] \\ 1, & x \in \left(\frac{1}{3}, \frac{2}{3}\right) \\ 0, & x \in \left[\frac{2}{3}, 1\right]. \end{cases} \quad (20)$$

$$u|_{\partial\Omega} = 0, \quad u := u(x) \in C^1(\Omega) \cap C(\bar{\Omega}), \quad x \in \Omega := (0, 1). \quad (21)$$

5 Solution of the BVP-2

The exact solution of the BVP given in (20) is

$$u'' = \begin{cases} -\frac{1}{6}x, & x \in \left[0, \frac{1}{3}\right]; \\ \frac{1}{2}\left(x^2 - x + \frac{1}{9}\right), & x \in \left(\frac{1}{3}, \frac{2}{3}\right); \\ \frac{1}{6}(x-1), & x \in \left[\frac{2}{3}, 1\right]. \end{cases} \quad (22)$$

The numerical solution of the BVP-2 is given both in numbers (see Table-2) and in graphical representation (see Figure-1). The result of this paper shows a regular fluctuating behavior with respect to both the finite difference and the exact results. As moving from two adjacent points one of the values approximates from the upper side and the other value from the lower side. This fluctuating behaviour gives us a possibility of offering better approximations. For example, it is clear from the figure and can be checked from the table that the arithmetical average of the successive function values almost coincides with the corresponding finite difference results. Further investigations are needed to develop new strategies to have better approximations.

Table-2 Numerical Results for the BVP-2

Figure-1 Graphical Representation of the Numerical Results for the BVP-2

Results for n=32	MIN % ERROR		MAX % ERROR		u1	u2	u3	u4	u5	u6	u7	u8
% error VASHA					5.00	2.50	9.98	4.37	11.02	4.99	11.44	5.33
u_VASHA	2.50	u2	11.91	u11	-0.0054688	-0.01016	-0.01719	-0.01992	-0.02891	-0.02969	-0.04063	-0.03945
u_EXACT					-0.0052083	-0.01042	-0.01563	-0.02083	-0.02604	-0.03125	-0.03646	-0.04167
u_DIRECT_FD	2.66	u16	3.21	u11	-0.0053711	-0.01074	-0.01611	-0.02148	-0.02686	-0.03223	-0.03760	-0.04297
% error D_FD					3.12	3.07	3.07	3.12	3.15	3.14	3.13	3.12
Results for n=32	u9	u10	u11	u12	u13	u14	u15	u16	<div>$u'' = \begin{cases} 0, & x \in [0, \frac{1}{2}]; \\ 1, & x \in (\frac{1}{2}, \frac{3}{2}); \\ 0, & x \in [\frac{3}{2}, 1]; \end{cases}$$u_0 = \begin{cases} \frac{1}{2}x, & x \in [0, \frac{1}{2}]; \\ \frac{1}{2}(x^2 - x + \frac{1}{2}), & x \in (\frac{1}{2}, \frac{3}{2}); \\ \frac{1}{2}(x-1), & x \in [\frac{3}{2}, 1]; \end{cases}$<p>$u _{\partial\Omega} = 0, u = u(x) \in C^1(\bar{\Omega}) \cap C(\bar{\Omega}), x \in \bar{\Omega} = (0,1).$</p><p>Note that the remaining values are symmetric with respect to value of u16.</p></div>			
% error VASHA	11.65	5.49	11.91	4.30	10.48	3.93	9.89	3.80				
u_VASHA	-0.05234	-0.04922	-0.06406	-0.05898	-0.07187	-0.06484	-0.07578	-0.06680				
u_EXACT	-0.04688	-0.05208	-0.05724	-0.06163	-0.06505	-0.06749	-0.06896	-0.06944				
u_DIRECT_FD	-0.04834	-0.05371	-0.05908	-0.06348	-0.06689	-0.06934	-0.07080	-0.07129				
% error D_FD	3.11	3.13	3.21	3.00	2.83	2.74	2.67	2.66				




Figure 1: Graphical Representation of the Numerical Results for the BVP-2

References

1. Cassel, K W. Variational Methods with Applications in Science and Engineering. Cambridge University Press, 2013.
2. Vashakmadze, T.S. Numerical Analysis I. TSU Press (in Georgian), 2009.